

Supplementary Table 2. Comparison of sarcopenia using grip strength

Participants	None	Sarcopenia	Total
Male			
60–64	178 (96.7)	6 (3.3)	184 (100.0)
65–69	203 (94.4)	12 (5.6)	215 (100.0)
70–74	141 (89.8)	16 (10.2)	157 (100.0)
75–79	105 (82.7)	22 (17.3)	127 (100.0)
≥80	52 (65.8)	27 (34.2)	79 (100.0)
Female			
60–64	262 (95.3)	13 (4.7)	275 (100.0)
65–69	219 (92.4)	18 (7.6)	237 (100.0)
70–74	152 (87.4)	22 (12.6)	174 (100.0)
75–79	108 (76.6)	33 (23.4)	141 (100.0)
≥80	53 (62.4)	32 (37.6)	85 (100.0)
Male			
≥60	679 (89.1)	83 (10.9)	762 (100.0)
≥65	501 (86.7)	77 (13.3)	578 (100.0)
≥70	298 (82.1)	65 (17.9)	363 (100.0)
≥75	157 (76.2)	49 (23.8)	206 (100.0)
≥80	52 (65.8)	27 (34.2)	79 (100.0)
Female			
≥60	794 (87.1)	118 (12.9)	912 (100.0)
≥65	532 (83.5)	105 (16.5)	637 (100.0)
≥70	313 (78.2)	87 (21.8)	400 (100.0)
≥75	161 (71.2)	65 (28.8)	226 (100.0)
≥80	53 (62.4)	32 (37.6)	85 (100.0)

The data is presented as N (%).